



CLASS 10 ACADEMIC PROGRAM-2020

HIGHER MATH

Lecture : HM-35

Chapter 11 : Coordinate Geometry




$$x = \sqrt{\frac{c^2}{2}} + c - \frac{b}{2}$$



উদ্ভাস

একাডেমিক এন্ড এডমিশন কেয়ার



www.udvash.com

SLOPE OF A LINE CONNECTING TWO POINTS

$$\text{Slope, } m = \frac{\text{change of Ordinate}}{\text{change of Abscissa}} = \frac{\Delta y}{\Delta x}$$


$$m_{AB} = \frac{7-3}{5-2} = \frac{4}{3}$$

$$m_{BA} = \frac{3-7}{2-5} = \frac{-4}{-3} = \frac{4}{3}$$

$$\text{Slope, } m = \frac{\text{rise}}{\text{run}}$$

$$= \frac{7-3}{5-2}$$

$$= \frac{4}{3}$$



- Find the slope of the line passing through the points $A(6, -1)$ and $B(3, 4)$. (Similar to Example: 17)

Figure needed $\rightarrow$ No

$$m_{AB} = \frac{-1 - 4}{6 - 3}$$

$$= \frac{-5}{3}$$


$$= -\frac{5}{3}$$

(Ans)

$$m = \frac{\text{rise}}{\text{run}}$$

$$m = \frac{\Delta y}{\Delta x}$$

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$



$BP = \text{rise}$

$AP = \text{run}$

POLL QUESTION-01

What is the slope of the line passing through the points $A(2, 7)$ এবং $B(-1, 13)$?

$x_2 \ y_2$

A	-6
B	-2
C	2
D	6

$$\begin{aligned} m_{AB} &= \frac{7-13}{2-(-1)} \\ &= \frac{-6}{3} \\ &= -2 \end{aligned}$$

$$m = \frac{\Delta y}{\Delta x}$$

$$\checkmark m = \frac{y_1 - y_2}{x_1 - x_2}$$


$$\checkmark m = \frac{y_2 - y_1}{x_2 - x_1}$$

SLOPE OF A STRAIGHT LINE

- The tangent (tan) of the inclination angle of a line with **positive x axis** is called the slope or gradient of a line (m).

$$m_{AB} = \tan(\angle ABX) \\ = \tan \theta$$

$$m_{CD} = \tan(\angle CDX) \\ = \tan(\pi - \alpha)$$



$$\tan \theta = \frac{AN}{BN} = \frac{\text{rise}}{\text{run}} = m$$

POLL QUESTION-02


The angle of inclination of a line is 135° to the negative x axis, what will be its slope?



$$m = \tan 45^\circ$$

A	$\tan 45$
B	$\tan 135$
C	Both A and B
D	Not possible

$$m = \tan \theta$$

θ = inclination
angle to positive
x axis






- The coordinates of the three points A, B and C are respectively $(2, 2)$, $(5, 2)$ and $(2, 7)$. Draw the lines AB and AC on the Cartesian plane. If possible, find the slopes of the lines AB and AC . (Example: 16)

$$m_{AB} = \frac{2-2}{5-2} = 0$$

$$m_{AB} = \tan(0) = 0$$

$$m_{AC} = \frac{7-2}{2-2} = \text{undefined}$$

$$m_{AC} = \tan(90) = \infty$$




(Ans)

POLL QUESTION-03

$$A(-2, 2), B(3, 2) \rightarrow m_{AB} = ?$$

$$m = \frac{2 - 2}{3 - (-2)} \\ = 0$$

A	1
B	-1
C	2
D	0



$$m = \tan(0) \\ = 0$$

- Find the value of p if the line joining the points $A(3, 3p)$ and $B(4, p^2 + 1)$ has slope -1 .
(Exercise: 5)

$$m_{AB} = \frac{p^2 + 1 - 3p}{4 - 3}$$


$$= \frac{p^2 - 3p + 1}{1}$$

A.T.Q. $p^2 - 3p + 1 = -1$

• • •

$$p = 1, 2$$

SLOPE OF COLLINEAR POINTS / PARALLEL LINES




slope equal

$$m_{AB} = \tan \theta$$

$$m_{BC} = \tan \theta$$

$$m_{AC} = \tan \theta$$



$$PQ \parallel RS$$

$$m_{PQ} = m_{RS}$$



উদ্ভাস

একাডেমিক এন্ড এডমিশন কন্ট্রোল

Higher Math


Chapter 11 : Coordinate Geometry

□ Show that the points $A(0, -3)$, $B(4, -2)$ and $C(16, 1)$ are collinear. (Exercise: 3)

$$m_{AB} = \frac{-2 - (-3)}{4 - 0} = \frac{1}{4}$$

$$(m_{AB} = m_{BC}) = m_{AC}$$

$$m_{BC} = \frac{1 - (-2)}{16 - 4} = \frac{3}{12} = \frac{1}{4}$$



∴ A, B, C are collinear.

- If the points $A(1, -1)$, $B(t, 2)$ and $C(t^2, t + 3)$ are collinear, find the admissible value of t .
(Exercise: 4)

$$m_{AC} = m_{AB}$$
$$\frac{t+3-(-1)}{t^2-1} = \frac{2-(-1)}{t-1}$$

or,

$$\frac{t+4}{t^2-1} = \frac{3}{t-1}$$

$$t = \frac{1}{2}, 1$$

$$t-1=0$$

POLL QUESTION-04

The slope of line PQ is 2 and RS is a line parallel to PQ. What is the slope of line RS?

RS || PQ

$$m_{RS} = m_{PQ}$$
$$= 2$$

A	$\frac{1}{2}$
B	$-\frac{1}{2}$
C	2
D	Not defined


- $A(t, 3t), B(t^2, 2t), C(t - 2, t)$ and $D(1, 1)$ are the four different points. If the lines AB and CD are parallel, find the admissible value of t . (Example: 19)

$$m_{AB} = m_{CD}$$

$$\frac{2t - 3t}{t^2 - t} = \frac{t - 1}{t - 2 - 1}$$

$$\frac{-t}{t(t-1)} = \frac{t-1}{t-3}$$

$$t = -1, 2$$



$$t \neq 0 \quad \text{How?}$$

□ Prove that the points $A(a, 0)$, $B(0, b)$ and $C(1, 1)$ are collinear if $\frac{1}{a} + \frac{1}{b} = 1$. (Exercise: 6)

Let, A, B, C be collinear points

$$m_{AC} = m_{BC}$$

$$\text{or, } \frac{0-1}{a-1} = \frac{b-1}{0-1}$$

$$\text{or, } \frac{-1}{a-1} = \frac{b-1}{-1}$$

$$\text{or, } (-1)(-1) = (b-1)(a-1)$$

$$\text{or, } 1 = ab - b - a + 1$$

$$\text{or, } b + a = ab$$

$$\text{or, } \frac{b+a}{ab} = \frac{ab}{ab}$$

$$\text{or, } \frac{b}{ab} + \frac{a}{ab} = 1$$

$$\therefore \frac{1}{a} + \frac{1}{b} = 1$$



উদ্ভাস

একাডেমিক এন্ড এডমিশন কেয়ার


Higher Math

Chapter 11 : Coordinate Geometry

Onnoro kom
path shala

Khan
Academy

Hand raise



না বুঝে
মুখস্থ করার
অভ্যাস প্রতিভাকে
ধ্বংস করে

$$X = \frac{c \rho V^2}{2S}$$

$$X = \frac{c \rho V^2}{2S}$$

$$E = mc^2$$

$$x = \sqrt{\frac{a^2}{c} + c} - \frac{b}{2}$$



উদ্ভাস

একাডেমিক এন্ড এডমিশন কোয়ার

www.udvash.com